

POLYNOMIALS

**It is not once nor twice but times without number that the same ideas
make their appearance in the world.**

1. Find the value for K for which $x^4 + 10x^3 + 25x^2 + 15x + K$ exactly divisible by $x + 7$.

(Ans : K= - 91)

Ans: Let $P(x) = x^4 + 10x^3 + 25x^2 + 15x + K$ and $g(x) = x + 7$

Since $P(x)$ exactly divisible by $g(x)$

$$\therefore r(x) = 0$$

now $x + 7 \overline{)x^4 + 10x^3 + 25x^2 + 15x + K}$
 $x^4 + 7x^3$

$$3x^3 + 25x^2$$

$$3x^3 + 21x^2$$

$$4x^2 + 15x$$

$$4x^2 + 28x$$

$$-13x + K$$

$$-13x - 91$$

$$K + 91$$

$$\therefore K + 91 = 0$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

K= -91

2. If two zeros of the polynomial $f(x) = x^4 - 6x^3 - 26x^2 + 138x - 35$ are $2 \pm \sqrt{3}$. Find the other zeros.

(Ans: 7, -5)

Ans: Let the two zeros are $2 + \sqrt{3}$ and $2 - \sqrt{3}$

$$\begin{aligned}\text{Sum of Zeros} &= 2 + \sqrt{3} + 2 - \sqrt{3} \\ &= 4\end{aligned}$$

$$\begin{aligned}\text{Product of Zeros} &= (2 + \sqrt{3})(2 - \sqrt{3}) \\ &= 4 - 3 \\ &= 1\end{aligned}$$

Quadratic polynomial is $x^2 - (\text{sum})x + \text{Product}$

$$\begin{array}{r} x^2 - 2x - 35 \\ x^2 - 4x + 1 \overline{)x^4 - 6x^3 - 26x^2 + 138x - 35} \\ \underline{x^4 - 4x^3 + x^2} \\ \hline -2x^3 - 27x^2 + 138x \\ -2x^3 + 8x^2 - 2x \\ \hline -35x^2 + 140x - 35 \\ -35x^2 + 140x - 35 \\ \hline 0 \end{array}$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

$$\therefore x^2 - 2x - 35 = 0$$

$$(x - 7)(x + 5) = 0$$

$$x = 7, -5$$

other two Zeros are 7 and -5

3. Find the Quadratic polynomial whose sum and product of zeros are $\sqrt{2} + 1$, $\frac{1}{\sqrt{2} + 1}$.

Ans: sum = $2\sqrt{2}$

Product = 1

Q.P =

$$X^2 - (\text{sum}) x + \text{Product}$$

$$\therefore x^2 - (2\sqrt{2})x + 1$$

4. If α, β are the zeros of the polynomial $2x^2 - 4x + 5$ find the value of a) $\alpha^2 + \beta^2$ b)
 $(\alpha - \beta)^2$

(Ans: a) -1 , b) -6)

Ans: p(x) = $2x^2 - 4x + 5$

$$\alpha + \beta = \frac{-b}{a} = \frac{4}{2} = 2$$

$$\alpha \beta = \frac{c}{a} = \frac{5}{2}$$

$$\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$$

$$\text{Substitute then we get, } \alpha^2 + \beta^2 = -1$$

$$(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

Substitute, we get $= (\alpha - \beta)^2 = -6$

5. If α, β are the zeros of the polynomial $x^2 + 8x + 6$ frame a Quadratic polynomial

whose zeros are a) $\frac{1}{\alpha}$ and $\frac{1}{\beta}$ b) $1 + \frac{\beta}{\alpha}, 1 + \frac{\alpha}{\beta}$.

(Ans: $x^2 + \frac{4}{3}x + \frac{1}{6}$, $x^2 - \frac{32}{3}x + \frac{32}{3}$)

Ans: $p(x) = x^2 + 8x + 6$

$$\alpha + \beta = -8 \text{ and } \alpha \beta = 6$$

a) Let two zeros are $\frac{1}{\alpha}$ and $\frac{1}{\beta}$

$$\text{Sum} = \frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha \cdot \beta} = \frac{-8}{6} = \frac{-4}{3}$$

$$\text{Product} = \frac{1}{\alpha} \times \frac{1}{\beta} = \frac{1}{\alpha \cdot \beta} = \frac{1}{6}$$

Required Q.P is

$$x^2 + \frac{4}{3}x + \frac{1}{6}$$

b) Let two Zeros are $1 + \frac{\beta}{\alpha}$ and $1 + \frac{\alpha}{\beta}$

$$\text{sum} = 1 + \frac{\beta}{\alpha} + 1 + \frac{\alpha}{\beta}$$

$$= 2 + \frac{\alpha}{\beta} + \frac{\beta}{\alpha}$$

$$= 2 + \frac{\alpha^2 + \beta^2}{\alpha \beta}$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

$$= 2 + \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} \text{ after solving this problem,}$$

We get $= \frac{32}{3}$

$$\text{Product} = \left(1 + \frac{\beta}{\alpha}\right)\left(1 + \frac{\alpha}{\beta}\right)$$

$$= 1 + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + 1$$

$$= 2 + \frac{\alpha^2 + \beta^2}{\alpha\beta}$$

Substitute this sum,

We get $= \frac{32}{3}$

$$\text{Required Q.P. is } x^2 - \frac{32}{3}x + \frac{32}{3}$$

6. On dividing the polynomial $4x^4 - 5x^3 - 39x^2 - 46x - 2$ by the polynomial $g(x)$ the quotient is $x^2 - 3x - 5$ and the remainder is $-5x + 8$. Find the polynomial $g(x)$.
 (Ans: $4x^2 + 7x + 2$)

Ans: $p(x) = g(x)q(x) + r(x)$

$$g(x) = \frac{p(x) - r(x)}{q(x)}$$

$$\text{let } p(x) = 4x^4 - 5x^3 - 39x^2 - 46x - 2$$

$$q(x) = x^2 - 3x - 5 \text{ and } r(x) = -5x + 8$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

$$\text{now } p(x) - r(x) = 4x^4 - 5x^3 - 39x^2 - 41x - 10$$

$$\text{when } \frac{p(x) - r(x)}{q(x)} = 4x^2 + 7x + 2$$

$$\therefore g(x) = 4x^2 + 7x + 2$$

7. If the squared difference of the zeros of the quadratic polynomial $x^2 + px + 45$ is equal to 144 , find the value of p. (Ans: ± 18).

Ans: Let two zeros are α and β where $\alpha > \beta$

According given condition

$$(\alpha - \beta)^2 = 144$$

$$\text{Let } p(x) = x^2 + px + 45$$

$$\alpha + \beta = \frac{-b}{a} = \frac{-p}{1} = -p$$

$$\alpha\beta = \frac{c}{a} = \frac{45}{1} = 45$$

$$\text{now } (\alpha - \beta)^2 = 144$$

$$(\alpha + \beta)^2 - 4\alpha\beta = 144$$

$$(-p)^2 - 4(45) = 144$$

Solving this we get $p = \pm 18$

8. If α, β are the zeros of a Quadratic polynomial such that $\alpha + \beta = 24$, $\alpha - \beta = 8$. Find a Quadratic polynomial having α and β as its zeros. (Ans: $k(x^2 - 24x + 128)$)

Ans: $\alpha + \beta = 24$

$$\alpha - \beta = 8$$

$$2\alpha = 32$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

$$\alpha = \frac{32}{2} = 16, \therefore \alpha = 16$$

Work the same way to $\alpha + \beta = 24$

$$\text{So, } \beta = 8$$

$$\begin{aligned} Q.P \text{ is } & x^2 - (\text{sum}) x + \text{product} \\ &= x^2 - (16+8) x + 16 \times 8 \end{aligned}$$

Solve this,

$$\text{it is } k(x^2 - 24x + 128)$$

9. If α & β are the zeroes of the polynomial $2x^2 - 4x + 5$, then find the value of

- a. $\alpha^2 + \beta^2$ b. $1/\alpha + 1/\beta$ c. $(\alpha - \beta)^2$ d. $1/\alpha^2 + 1/\beta^2$ e. $\alpha^3 + \beta^3$

$$(\text{Ans:}-1, \frac{4}{5}, -6, \frac{-4}{25}, -7)$$

Ans: Let $p(x) = 2x^2 - 4x + 5$

$$\alpha + \beta = \frac{-b}{a} = \frac{4}{2} = 2$$

$$\alpha\beta = \frac{c}{a} = \frac{5}{2}$$

$$\text{a)} \quad \alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$$

$$\text{Substitute to get } \alpha^2 + \beta^2 = -1$$

$$\text{b)} \quad \frac{1}{a} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta}$$

$$\text{substitute, then we get } \frac{1}{a} + \frac{1}{\beta} = \frac{4}{5}$$

$$\text{b)} \quad (\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$$

$$\text{Therefore we get, } (\alpha - \beta)^2 = -6$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

$$d) \frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{\alpha^2 + \beta^2}{\alpha\beta^2} = \frac{-1}{\left(\frac{5}{2}\right)^2}$$

$$\therefore \frac{1}{\alpha^2} + \frac{1}{\beta^2} = \frac{-4}{25}$$

$$e) \alpha^3 + \beta^3 = (\alpha + \beta)(\alpha^2 + \beta^2 - \alpha\beta)$$

Substitute this,

to get, $\alpha^3 + \beta^3 = -7$

10. Obtain all the zeros of the polynomial $p(x) = 3x^4 - 15x^3 + 17x^2 + 5x - 6$ if two zeroes are $-1/\sqrt{3}$ and $1/\sqrt{3}$. (Ans:3,2)

11. Give examples of polynomials $p(x)$, $g(x)$, $q(x)$ and $r(x)$ which satisfy the division algorithm.

a. $\deg p(x) = \deg q(x)$ b. $\deg q(x) = \deg r(x)$ c. $\deg q(x) = 0$.

12. If the ratios of the polynomial $ax^3 + 3bx^2 + 3cx + d$ are in AP, Prove that $2b^3 - 3abc + a^2d = 0$

Ans: Let $p(x) = ax^3 + 3bx^2 + 3cx + d$ and α, β, γ are their three Zeros

but zero are in AP

let $\alpha = m - n$, $\beta = m$, $\gamma = m + n$

$$\text{sum} = \alpha + \beta + \gamma = \frac{-b}{a}$$

$$\text{substitute this sum, to get } m = \frac{-b}{a}$$

UNIT-2 POLYNOMIALS

Downloaded From: www.jsuniltutorial.weebly.com

[Year]

Now taking two zeros as sum $\alpha\beta + \beta r + \alpha r = \frac{c}{a}$

$$(m-n)m + m(m+n) + (m+n)(m-n) = \frac{3c}{a}$$

Solve this problem , then we get

$$\frac{3b^2 - 3ac}{a^2} = n^2$$

$$\text{Product } \alpha\beta r = \frac{d}{a}$$

$$(m-n)m (m+n) = \frac{-d}{a}$$


$$(m^2 - n^2)m = \frac{-d}{a}$$

$$[(\frac{-b}{a})^2 - (\frac{3b^2 - 3ac}{a^2})] (\frac{-b}{a}) = \frac{-d}{a}$$

Simplifying we get

$$2b^3 - 3abc + a^2 d = 0$$

13. Find the number of zeros of the polynomial from the graph given.



(Ans:1)

14. If one zero of the polynomial $3x^2 - 8x + 2k+1$ is seven times the other, find the zeros and the value of k
 (Ans k= 2/3)

Self Practice

14. If $(n-k)$ is a factor of the polynomials x^2+px+q & $x^2 + m x+n$. Prove that

$$k = n + \frac{n - q}{m - p}$$

Ans : since $(n - k)$ is a factor of $x^2 + px + q$

$$\therefore (n - k)^2 + p(n - k) + q = 0$$

$$\text{And } (n - k)^2 + m(n - k) + n = 0$$

Solve this problem by yourself,

$$\therefore k = n + \frac{n - q}{m - p}$$

SELF PRACTICE

16. If $2, \frac{1}{2}$ are the zeros of px^2+5x+r , prove that $p= r$.

17. If m, n are zeroes of ax^2-5x+c , find the value of a and c if $m + n = m.n=10$

(Ans: a=1/2 ,c=5)

18. What must be subtracted from $8x^4 + 14x^3 - 2x^2 + 7x - 8$ so that the resulting polynomial is exactly divisible by $4x^2+3x-2$.
 (Ans: $14x - 10$)

19. What must be added to the polynomial $p(x)=x^4 + 2x^3 - 2x^2 + x - 1$ so that the resulting polynomial is exactly divisible by x^2+2x-3 .
 (Ans: x-2)