UNIT-1

#### **NUMBER SYSTEMS**

Numbers are intellectual witnesses that belong only to mankind.

1. If the H C F of 657 and 963 is expressible in the form of 657x + 963x - 15 find x. (Ans:x=22)

**Ans:** Using Euclid's Division Lemma

$$a=bq+r, o \le r < b$$

$$963=657\times1+306$$

$$657=306\times2+45$$

$$306=45\times6+36$$

$$45=36\times1+9$$

$$36=9\times4+0$$
∴ HCF (657, 963) = 9
$$now \quad 9 = 657x + 963\times(-15)$$

$$657x=9+963\times15$$

$$=9+14445$$

$$657x=14454$$

$$x=14454/657$$

x = 22

2. Express the GCD of 48 and 18 as a linear combination. (Ans: Not unique)

A=bq+r, where 
$$o \le r < b$$
  
 $48=18x2+12$   
 $18=12x1+6$   
 $12=6x2+0$   
 $\therefore$  HCF  $(18,48) = 6$   
now  $6=18-12x1$   
 $6=18-(48-18x2)$   
 $6=18-48x1+18x2$   
 $6=18x3-48x1$   
 $6=18x3+48x(-1)$   
 $6=18x+48y$   
 $x=3, y=-1$ 

i.e.

*:*.

### JSUNIL TUTORIAL ACBSE Coaching for Mathematics and Science

$$x = 51, y = -19$$

Hence, x and y are not unique.

3. Prove that one of every three consecutive integers is divisible by 3.

#### Ans:

n,n+1,n+2 be three consecutive positive integers We know that n is of the form 3q, 3q + 1, 3q + 2So we have the following cases

Case – I when 
$$n = 3q$$

In the this case, n is divisible by 3 but n + 1 and n + 2 are not divisible by 3

Case - II When 
$$n = 3q + 1$$
  
Sub  $n = 2 = 3q + 1 + 2 = 3(q + 1)$  is divisible by 3. but n and n+1 are not divisible by 3

Case – III When 
$$n = 3q + 2$$
  
Sub  $n = 2 = 3q + 1 + 2 = 3(q + 1)$  is divisible by 3. but n and n+1 are not divisible by 3

Hence one of n, n + 1 and n + 2 is divisible by 3

4. Find the largest possible positive integer that will divide 398, 436, and 542 leaving remainder 7, 11, 15 respectively.

(Ans: 17)

Ans: The required number is the HCF of the numbers

Find the HCF of 391, 425 and 527 by Euclid's algorithm

$$\therefore$$
 HCF (425, 391) = 17

Now we have to find the HCF of 17 and 527  $527 = 17 \times 31 + 0$ 

## JSUNIL TUTORIAL ACBSE Coaching for Mathematics and Science

$$\therefore$$
 HCF (17,527) = 17  
  $\therefore$  HCF (391, 425 and 527) = 17

5. Find the least number that is divisible by all numbers between 1 and 10 (both inclusive).

(Ans:2520)

**Ans:** The required number is the LCM of 1,2,3,4,5,6,7,8,9,10

$$\therefore$$
 LCM = 2  $\times$  2  $\times$  3  $\times$  2  $\times$  3  $\times$  5  $\times$  7 = 2520

6. Show that 571 is a prime number.

Ans: Let  $x=571 \Rightarrow \sqrt{x}=\sqrt{571}$ 

Now 571 lies between the perfect squares of  $(23)^2$  and  $(24)^2$  Prime numbers less than 24 are 2,3,5,7,11,13,17,19,23 Since 571 is not divisible by any of the above numbers 571 is a prime number

7. If d is the HCF of 30, 72, find the value of x & y satisfying d = 30x + 72y.

(Ans:5, -2 (Not unique)

**Ans:** Using Euclid's algorithm, the HCF (30, 72)

$$72 = 30 \times 2 + 12$$

$$30 = 12 \times 2 + 6$$

$$12 = 6 \times 2 + 0$$

$$HCF(30,72) = 6$$

$$6=30-(72-30\times2)2$$

$$6=30-2\times72+30\times4$$

$$6=30\times5+72\times-2$$

$$\therefore x = 5, y = -2$$

Also 
$$6 = 30 \times 5 + 72 (-2) + 30 \times 72 - 30 \times 72$$

Solve it, to get

$$x = 77, y = -32$$

Hence, x and y are not unique

# JSUNIL TUTORIAL ACBSE Coaching for Mathematics and Science

8. Show that the product of 3 consecutive positive integers is divisible by 6.

Ans: Proceed as in question sum no. 3

9. Show that for odd positive integer to be a perfect square, it should be of the form 8k +1.

Let a=2m+1

Ans: Squaring both sides we get

$$a^2 = 4m (m + 1) + 1$$

.. product of two consecutive numbers is always even

$$m(m+1)=2k$$

$$a^2 = 4(2k) + 1$$

$$a^2 = 8 k + 1$$

Hence proved

10. Find the greatest number of 6 digits exactly divisible by 24, 15 and 36. (Ans:999720)

Ans: LCM of 24, 15, 36

$$LCM = 3 \times 2 \times 2 \times 2 \times 3 \times 5 = 360$$

Now, the greatest six digit number is 999999

$$\therefore$$
 the required number = 999999 - 279 = 999720

11. If a and b are positive integers. Show that  $\sqrt{2}$  always lies between  $\frac{a}{b}$  and  $\frac{a-2b}{a+b}$ 

 $\therefore$  Q = 2777, R = 279

Ans: We do not know whether  $\frac{a^2 - 2b^2}{b(a+b)}$  or  $\frac{a}{b} < \frac{a+2b}{a+b}$ 

:. to compare these two number,

Let us comute  $\frac{a}{b} - \frac{a+2b}{a+b}$ 

=> on simplifying , we get 
$$\frac{a^2 - 2b^2}{b(a+b)}$$

### JSUNIL TUTORIAL

ACBSE Coaching for Mathematics and Science

$$\therefore \frac{a}{b} - \frac{a+2b}{a+b} > 0 \text{ or } \frac{a}{b} - \frac{a+2b}{a+b} < 0$$

$$now \frac{a}{b} - \frac{a+2b}{a+b} > 0$$

$$\frac{a^2 - 2b^2}{b(a+b)} > 0 \quad \text{solve it, we get, } a > \sqrt{2b}$$

Thus , when a > 
$$\sqrt{2}b$$
 and  $\frac{a}{b} < \frac{a+2b}{a+b}$ ,

We have to prove that 
$$\frac{a+2b}{a+b} < \sqrt{2} < \frac{a}{b}$$

Now a 
$$>\sqrt{2}$$
 b $\Rightarrow$ 2a<sup>2</sup>+2b<sup>2</sup>>2b<sup>2</sup>+ a<sup>2</sup>+2b<sup>2</sup>  
On simplifying we get

$$\sqrt{2} > \frac{a+2b}{a+b}$$

Also a>
$$\sqrt{2}$$

$$\Rightarrow \frac{a}{b} > \sqrt{2}$$

Similarly we get 
$$\sqrt{2}$$
,  $<\frac{a+2b}{a+b}$ 

Hence 
$$\frac{a}{b} < \sqrt{2} < \frac{a+2b}{a+b}$$

12. Prove that 
$$(\sqrt{n-1} + \sqrt{n+1})$$
 is irrational, for every  $n \in \mathbb{N}$ 

#### **Self Practice**