# 10th Chapter Number System CBSE Test Paper - 03

BSE Coaching for Mathematics and S

Numbers are intellectual witnesses that belong only to mankind.

- 1. If the H C F of 657 and 963 is expressible in the form of 657x + 963x 15 find x. (Ans:x=22)
- **Ans:** Using Euclid's Division Lemma

```
a = bq+r, o \le r < b
963=657 \times 1+306
657=306 \times 2+45
306=45 \times 6+36
45=36 \times 1+9
36=9 \times 4+0
\therefore \text{ HCF } (657, 963) = 9
now \quad 9 = 657x + 963 \times (-15)
657x=9+963 \times 15
=9+14445
657x=14454
x=14454/657
x = 22
```

2. Express the GCD of 48 and 18 as a linear combination. (Ans: Not unique)

```
A=bq+r, where o \le r < b

48=18x2+12

18=12x1+6

12=6x2+0

\therefore HCF (18,48) = 6

now 6= 18-12x1

6= 18-(48-18x2)

6= 18-48x1+18x2

6= 18x3-48x1

6= 18x3+48x(-1)

6= 18x +48y

x=3, y=-1
```

i.e.

*.*..



*.*..

Hence, x and y are not unique.

x = 51, y = -19

3. Prove that one of every three consecutive integers is divisible by 3.

#### Ans:

n,n+1,n+2 be three consecutive positive integers We know that n is of the form 3q, 3q + 1, 3q + 2So we have the following cases

Case -I when n = 3q

In the this case, n is divisible by 3 but n + 1 and n + 2 are not divisible by 3

Case - II When n = 3q + 1Sub n = 2 = 3q + 1 + 2 = 3(q + 1) is divisible by 3. but n and n+1 are not divisible by 3

Case – III When n = 3q + 2Sub n = 2 = 3q + 1 + 2 = 3(q + 1) is divisible by 3. but n and n+1 are not divisible by 3

Hence one of n, n + 1 and n + 2 is divisible by 3

4. Find the largest possible positive integer that will divide 398, 436, and 542 leaving remainder 7, 11, 15 respectively.
 (Ans: 17)

Ans: The required number is the HCF of the numbers

Find the HCF of 391, 425 and 527 by Euclid's algorithm

: HCF (425, 391) = 17

Now we have to find the HCF of 17 and 527 527 = 17 x 31 +0



∴ HCF (17,527) = 17 ∴ HCF (391, 425 and 527) = 17

5. Find the least number that is divisible by all numbers between 1 and 10 (both inclusive).

(Ans:2520)

**Ans:** The required number is the LCM of 1,2,3,4,5,6,7,8,9,10

 $\therefore \text{ LCM} = 2 \times 2 \times 3 \times 2 \times 3 \times 5 \times 7 = 2520$ 

6. Show that 571 is a prime number.

Ans: Let  $x=571 \Rightarrow \sqrt{x}=\sqrt{571}$ 

Now 571 lies between the perfect squares of  $(23)^2$  and  $(24)^2$ Prime numbers less than 24 are 2,3,5,7,11,13,17,19,23 Since 571 is not divisible by any of the above numbers 571 is a prime number

7. If d is the HCF of 30, 72, find the value of x & y satisfying d = 30x + 72y. (Ans:5, -2 (Not unique)

**Ans:** Using Euclid's algorithm, the HCF (30, 72)

Also  $6 = 30 \times 5 + 72 (-2) + 30 \times 72 - 30 \times 72$ 

Solve it, to get

$$x = 77, y = -32$$

Hence, x and y are not unique

#### www.jsuniltutorial.weebly.com/

#### page - 03

#### 8. Show that the product of 3 consecutive positive integers is divisible by 6.

Ans: Proceed as in question sum no. 3

9. Show that for odd positive integer to be a perfect square, it should be of the form 8k + 1.

BSE Coaching for Mathematics and Science

Let a=2m+1

Ans: Squaring both sides we get

$$a^2 = 4m(m+1) + 1$$

 $\therefore$  product of two consecutive numbers is always even

m(m+1)=2k $a^{2}=4(2k)+1$  $a^{2} = 8 k + 1$ Hence proved

10. Find the greatest number of 6 digits exactly divisible by 24, 15 and 36. (Ans:999720)

Ans: LCM of 24, 15, 36

 $LCM = 3 \times 2 \times 2 \times 2 \times 3 \times 5 = 360$ 

Now, the greatest six digit number is 999999 Divide 999999 by 360  $\therefore Q = 2777$ , R = 279

: the required number = 999999 - 279 = 999720

11. If a and b are positive integers. Show that  $\sqrt{2}$  always lies between  $\frac{a}{b}$  and  $\frac{a-2b}{a+b}$ 

**Ans:** We do not know whether  $\frac{a^2 - 2b^2}{b(a+b)}$  or  $\frac{a}{b} < \frac{a+2b}{a+b}$ 

 $\therefore$  to compare these two number,

Let us comute 
$$\frac{a}{b} - \frac{a+2b}{a+b}$$
  
=> on simplifying , we get  $\frac{a^2 - 2b^2}{b(a+b)}$ 

# BSE Coaching for Mathematics and Science

$$\therefore \frac{a}{b} - \frac{a+2b}{a+b} > 0 \text{ or } \frac{a}{b} - \frac{a+2b}{a+b} < 0$$

now 
$$\frac{a}{b} - \frac{a+2b}{a+b} > 0$$
  
 $\frac{a^2 - 2b^2}{b(a+b)} > 0$  solve it, we get,  $a > \sqrt{2b}$ 

$$(+b)$$

Thus , when a >  $\sqrt{2b}$  and  $\frac{a}{b} < \frac{a+2b}{a+b},$ 

We have to prove that  $\frac{a+2b}{a+b} < \sqrt{2} < \frac{a}{b}$ 

Now a  $>\sqrt{2} b \Longrightarrow 2a^2 + 2b^2 > 2b^2 + a^2 + 2b^2$ On simplifying we get

$$\sqrt{2} > \frac{a+2b}{a+b}$$
Also  $a > \sqrt{2}$ 

$$\Rightarrow \frac{a}{b} > \sqrt{2}$$
Similarly we get  $\sqrt{2}$ ,  $<\frac{a+2b}{a+b}$ 
Hence  $\frac{a}{b} < \sqrt{2} < \frac{a+2b}{a+b}$ 

12. Prove that  $(\sqrt{n-1} + \sqrt{n+1})$  is irrational, for every  $n \in \mathbb{N}$ 

#### **Self Practice**