**UNIT-1** 

## **NUMBER SYSTEMS**

Numbers are intellectual witnesses that belong only to mankind.

963x 1. If the H C F of 657 and 963 is expressible in the form of 657x + 963x - 15 find x. (Ans:x=22)

Ans: Using Euclid's Division Lemma a = bq + r,  $o \le r < b$ 

:. HCF (657, 963) = 9  
now 
$$9 = 657x + 963 \times (-15)$$

$$657x = 9 + 963 \times 15$$

$$= 9 + 14445$$

2. Express the GCD of 48 and 18 as a linear combination. (Ans: Not unique)

A=bq+r, where 
$$o \le r < b$$

$$48 = 18x2 + 12$$

$$18=12x1+6$$

$$12 = 6x2 + 0$$

$$\therefore$$
 HCF (18,48) = 6

now 
$$6 = 18 - 12x1$$

$$6 = 18 - (48 - 18x2)$$

$$6 = 18 - 48x1 + 18x2$$

$$6 = 18x3-48x1$$

$$6 = 18x3 + 48x(-1)$$

i.e. 
$$6 = 18x + 48y$$

$$x=3, y=-1$$

$$x = 51, y = -19$$

3. Prove that one of every three consecutive integers is divisible by 3.

Ans:

n,n+1 n+2 hord.

n,n+1,n+2 be three consecutive positive integers We know that n is of the form 3q, 3q + 1, 3q + 2So we have the following cases

Case 
$$-I$$
 when  $n = 3q$ 

In the this case, n is divisible by 3 but n+1 and n+2 are not divisible by 3

Case - II When 
$$n = 3q + 1$$
  
Sub  $n = 2 = 3q + 1 + 2 = 3(q + 1)$  is divisible by 3. but n and n+1 are not divisible by 3

Case – III When 
$$n = 3q + 2$$
  
Sub  $n = 2 = 3q + 1 + 2 = 3(q + 1)$  is divisible by 3. but n and n+1 are not divisible by 3

Hence one of n, n + 1 and n + 2 is divisible by 3

4. Find the largest possible positive integer that will divide 398, 436, and 542 leaving • remainder 7, 11, 15 respectively. (Ans: 17)

Ans: The required number is the HCF of the numbers

Find the HCF of 391, 425 and 527 by Euclid's algorithm

$$\therefore$$
 HCF (425, 391) = 17

Now we have to find the HCF of 17 and 527  $527 = 17 \times 31 + 0$ 

$$\therefore$$
 HCF (17,527) = 17  
  $\therefore$  HCF (391, 425 and 527) = 17

5. Find the least number that is divisible by all numbers between 1 and 10 (both inclusive).

(Ans:2520)

**Ans:** The required number is the LCM of 1,2,3,4,5,6,7,8,9,10

$$\therefore$$
 LCM = 2  $\times$  2  $\times$  3  $\times$  2  $\times$  3  $\times$  5  $\times$  7 = 2520

6. Show that 571 is a prime number.

Ans: Let  $x=571 \Rightarrow \sqrt{x} = \sqrt{571}$ 

Now 571 lies between the perfect squares of (23)<sup>2</sup> and (24) Prime numbers less than 24 are 2,3,5,7,11,13,17,19,23 Since 571 is not divisible by any of the above numbers 571 is a prime number



**Ans:** Using Euclid's algorithm, the HCF (30, 72)

$$72 = 30 \times 2 + 12$$
  
 $30 = 12 \times 2 + 6$   
 $12 = 6 \times 2 + 0$ 

HCF 
$$(30,72) \neq 6$$
  
 $6=30-12\times 2$   
 $6=30-(72-30\times 2)2$   
 $6=30-2\times 72+30\times 4$   
 $6=30\times 5+72\times -2$   
 $\therefore x = 5, y = -2$ 

Also 
$$6 = 30 \times 5 + 72 (-2) + 30 \times 72 - 30 \times 72$$

Solve it, to get

$$x = 77, y = -32$$

Hence, x and y are not unique

8. Show that the product of 3 consecutive positive integers is divisible by 6.

Ans: Proceed as in question sum no. 3

9. Show that for odd positive integer to be a perfect square, it should be of the form 8k + 1.

Let a=2m+1

Ans: Squaring both sides we get

$$a^2 = 4m (m + 1) + 1$$

... product of two consecutive numbers is always even

$$m(m+1)=2k$$

$$a^2 = 4(2k) + 1$$

$$a^2 = 8 k + 1$$

Hence proved

10. Find the greatest number of 6 digits exactly divisible by 24, 15 and 36. (Ans:999720)

Ans: LCM of 24, 15, 36

$$LCM = 3 \times 2 \times 2 \times 2 \times 3 \times 5 = 360$$

Now, the greatest six digit number is 999999

$$Q = 2777$$
,  $R = 279$ 

- $\therefore$  the required number = 999999 279 = 999720
- 11. If a and b are positive integers. Show that  $\sqrt{2}$  always lies between  $\frac{a}{b}$  and  $\frac{a-2b}{a+b}$

Ans: We do not know whether  $\frac{a^2 - 2b^2}{b(a+b)}$  or  $\frac{a}{b} < \frac{a+2b}{a+b}$ 

∴ to compare these two number,

Let us comute  $\frac{a}{b} - \frac{a+2b}{a+b}$ 

$$\Rightarrow$$
 on simplifying, we get  $\frac{a^2 - 2b^2}{b(a+b)}$ 

$$\therefore \frac{a}{b} - \frac{a+2b}{a+b} > 0 \text{ or } \frac{a}{b} - \frac{a+2b}{a+b} < 0$$

$$now \frac{a}{b} - \frac{a+2b}{a+b} > 0$$

$$\frac{a^2 - 2b^2}{b(a+b)} > 0 \quad \text{solve it , we get , } a > \sqrt{2b}$$

Thus , when 
$$a > \sqrt{2b}$$
 and  $\frac{a}{\sqrt{2b}} < \frac{a+2b}{\sqrt{2b}}$ ,

 $+ a^{2}+2b^{2}$   $\Rightarrow \frac{a}{b} > \sqrt{2}$ Similarly we get  $\sqrt{2}$ ,  $<\frac{a+2b}{a+b}$ Hence  $\frac{a}{b} < \sqrt{2} < \frac{a+2b}{a+b}$ 2. Prove that  $(\sqrt{p})$ 

$$\sqrt{2} > \frac{a + 2b}{a + b}$$

$$\Rightarrow \frac{a}{b} > \sqrt{2}$$

Hence 
$$\frac{a}{b} < \sqrt{2} < \frac{a+2b}{a+b}$$